US ERA ARCHIVE DOCUMENT

Mary Ann Rozum, USDA mrozum@csrees.usda.gov 202-401-4533

CSREES Sponsored Research on hormones

- 60 projects in USDA's Current Research Information Service database were identified as hormone-related
- Of the 60 research projects, 16 assessed the role of hormones in aquaculture and animal reproduction
- 44 projects investigated hormones in animal production and subsequent transport to the environment.
- Of the 44 projects, hormone-related research was a major component in 14 projects and a minor component in 30 projects.

Key Research Findings

- USDA ARS found that composting is an effective method to reduce, but not eliminate, the introduction of estradiol and testosterone into the environment
- Tennessee researchers comparing dairy and swine wastes concluded that swine farrowing facilities pose a greater threat as environmental estrogen pollutants.
- Purdue University studies determined that androgen and trenbolone and the antibiotics tylosin, monension, and lasalocid are highly sorbed to soils, have a relatively short half-life, thus concentrations in nearby water bodies are likely to be small.

Other Research Projects – minor focus on hormones

- Mississippi, Missouri, and Nebraska are researching hormone impacts in broiler, swine and cattle production.
- North Dakota State University is working with multi-state partners to study the fate and transport of estradiol and testosterone.
- West Virginia State University is investigating the fate of hormones during anaerobic digestion.
- Numerous other projects are investigating integrated manure management, animal production and soil flow transport processes.

Other Research Projects – minor focus on hormones

- ✓ ARS in Arkansas is investigating use of alum to decrease estrogen in runoff and forage-based livestock
- ✓ ARS in Georgia is investigating pathogen and hormone transport and researching whole farm management of agricultural effluent
- ✓ Iowa State University is investigating metal, pesticide and estrogen transport
- University of Maryland is investigating endocrine activity in poultry litter and subsequent impacts on water quality and fish and amphibian populations.

Key Research Findings (continued)

- Maryland scientists found that substantial quantities of poultry litter-derived estradiol can be transported to surface waters via runoff and persist for weeks to months at environmentally relevant concentrations.
- Also, fields under no-till management practices can lose up to ten times more estradiol than fields employing conventional tillage.